'O' PectiLyte™

Nutritional / Microbial Supplement for Calves

Consider for organic use

Contains a source of live (viable) naturally occurring microorganisms.

GUARANTEED ANALYSIS (per pound)

Crude Protein (min): 1 %
Crude Fiber (max): 6 %
Sodium (min): 9.2 %
Sodium (max): 11.0 %
Potassium (min): 2.6 %

Live Microbial Count:

40 billion colony forming units

(Lactobacillus acidophilus, Lactobacillus casei, Enterococcus faecium, Lactobacillus lactis, Pediococcus acidilactici, Bifidobacterium bifidum, Bifidobacterium infantis, Bifidobacterium longum, Bacillus licheniformis, Bacillus subtilis)

INGREDIENTS: Reed-sedge peat, sodium bicarbonate, sodium chloride, organic dried kelp, hydrolyzed yeast, potassium chloride, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, thiamine mononitrate, riboflavin, niacin, calcium pantothenate, pyridoxine hydrochloride, folic acid, biotin, vitamin B12 supplement, ascorbic acid, dried Lactobacillus acidophilus fermentation product, dried Lactobacillus casei fermentation product, dried Enterococcus faecium fermentation product, dried Lactobacillus lactis fermentation product, dried Pediococcus acidilactici fermentation product, dried Bifidobacterium bifidum fermentation product, dried Bifidobacterium infantis fermentation product, dried Bifidobacterium longum fermentation product, dried Bacillus subtilis fermentation product, dried Bacillus licheniformis fermentation product.

NOTICE

All microbial and enzyme sources in Bio-Vet products are naturally occurring (non-GMO). No Bio-Vet products are irradiated nor are any components derived from sewage sludge.

RECOMMENDED USES

Nutritional supplement for calves at shipping, during fluid loss, and when diarrhea is common among calves. Supplies dietary fibers, electrolytes and beneficial microbial cultures.

FEEDING DIRECTIONS

(1 scoop = 1 ounce)

For calves with fluid deficiency, continue feeding milk at alternate times (at least 30 minutes after feeding 'O' PectiLyte with water). Consult veterinarian for proper drenching/tubing procedures.

Daily maintenance: Mix 1/2 to 1 ounce 'O' PectiLyte / 2 quarts milk replacer or warm water. Feed once or twice daily.

At receiving: Mix 1/2 to 1 ounce 'O' PectiLyte / 2 quarts warm water once or twice daily at alternate times to milk.

Slight fluid deficiency: Mix 1 to 2 ounces 'O' PectiLyte / 2 quarts warm water twice daily. Moderate fluid deficiency: Mix 2 ounces 'O' PectiLyte / 2 quarts warm water two to three times daily.

Storage: Keep closed between uses. Store in cool, dry place.

PRODUCT CODE

OG-01520 11 pound pouch, 4 pouches / case



TS OPectiLyteRevised02092021

Bio-Vet, Inc. • 300 Ernie Drive • Barneveld, WI 53507, U.S.A. • (800) 246-8381 • Fax (608) 924-7006 • www.Bio-Vet.com